

String theory in Lorentz-invariant light cone gauge - II

Igor Nikitin, Lialia Nikitina

Fraunhofer Society, IMK.VE, 53754, St.Augustin, Germany

Abstract

We perform a quantization of 4-dimensional Nambu-Goto theory of open string in light cone gauge, related in Lorentz-invariant way with the world sheet. This allows to obtain a quantum theory without anomalies in Lorentz group. We consider a special type of gauge fixing conditions, imposed in oscillator sector of the theory, which lead to a relatively simple Hamiltonian mechanics, convenient for canonical quantization. We discuss the algebraic and geometric properties of this mechanics and determine its mass spectrum for the states of spin singlet $S = 0$.

1 Introduction

This paper continues the work [1], further referred as Part I, devoted to Dirac's quantization of Nambu-Goto theory of open string, formulated in the space-time of dimension $d=4$. The approach consists in the introduction of light cone gauge with the gauge axis related to a certain dynamical light-like vector in the theory. In this approach Lorentz group transforms the world sheet of the string together with the gauge axis and is not followed by reparametrizations. It is shown in Part I, that under these conditions the quantum Lorentz group is free of anomalies. The details of the mechanics depend on the choice of the dynamical vector to which the gauge axis is attached, i.e. on the choice of *gauge fixing conditions*. Generally the mechanics appears to be algebraically complex. In this paper, Section 2 we consider gauge fixing conditions, producing a resolvable system of polynomial equations. We discuss the structure of the obtained solutions, including their remaining discrete gauge symmetries. In Section 3 we perform quantization of this mechanics and determination of its mass spectrum for the states of spin singlet $S = 0$. In appendices we present the formulae for the solutions of polynomial system, provide proofs of the statements formulated throughout the paper and give the details about numerical methods used for determination of mass spectrum.

2 Classical mechanics

2.1 Lorentz-invariant abelian light cone gauge (lia-lcg)

We start from the canonical basis of string mechanics constructed in Part I:

$$P_\mu, Z_\mu + \text{infinite set of oscillators } a_k, a_k^* + \text{the top } \vec{e}_i, \vec{S},$$

with Poisson brackets

$$\begin{aligned} \{Z_\mu, P_\nu\} &= g_{\mu\nu}, \\ \{a_k, a_n^*\} &= ik\delta_{kn}, \quad k, n \in \mathbf{Z}/\{0\}, \\ \{S^i, S^j\} &= -\epsilon^{ijk}S^k, \quad \{S^i, e_n^j\} = -\epsilon^{ijk}e_n^k, \end{aligned} \quad (1)$$

and equivalent symplectic form

$$\Omega = dP_\mu \wedge dZ_\mu + \sum_{k \neq 0} \frac{1}{ik} da_k^* \wedge da_k + \frac{1}{2} d\vec{e}_i \wedge d(\vec{S} \times \vec{e}_i). \quad (2)$$

The mechanics is restricted by 4 constraints of the 1st class:

$$\{\chi_0, \chi_i\} = 0, \quad \{\chi_i, \chi_j\} = \epsilon_{ijk}\chi_k,$$

which include mass shell condition and requirements of the form “spin of the top is equal to the spin of the string”¹:

$$\begin{aligned} \chi_0 &= \frac{P^2}{2\pi} - L_0 = 0, \quad L_0 = \sum_{n \neq 0} a_n^* a_n, \\ \chi_3 &= S_3 - A_3 = 0, \quad A_3 = \sum_{n \neq 0} \frac{1}{n} a_n^* a_n, \\ \chi_+ &= S_+ - A_+ = 0, \quad \chi_- = S_- - A_- = 0, \\ \chi_\pm &= \chi_1 \pm i\chi_2, \quad S_\pm = S_1 \pm iS_2, \\ A_- &= \sqrt{\frac{2\pi}{P^2}} \sum_{k, n, k+n \neq 0} \frac{1}{k} a_k a_n a_{k+n}^*, \\ A_+ &= \sqrt{\frac{2\pi}{P^2}} \sum_{k, n, k+n \neq 0} \frac{1}{k} a_k^* a_n^* a_{k+n}, \end{aligned} \quad (3)$$

where $S_i = S^k e_i^k$ is a projection of $\vec{S}$ onto $\vec{e}_i$. Constraints generate the following transformations:

- χ_0 generates phase shifts of oscillator variables $E_0 : a_n \rightarrow a_n e^{-in\tau}$ and translations of mean coordinate $Z_\mu \rightarrow Z_\mu + P_\mu \tau / \pi$;
- χ_3 generates phase shifts $R_3 : a_n \rightarrow a_n e^{-i\alpha}$ and rotations of $\vec{e}_{1,2}$ about $\vec{e}_3$: $\vec{e}_i \rightarrow R(\vec{e}_3, -\alpha)\vec{e}_i$, $i = 1, 2$;
- $\chi_{1,2}$ generate rotations of basis $\vec{e}_i$ about axes $\vec{e}_{1,2}$ and certain non-linear transformations of oscillator variables, which will be considered in more details in subsection 2.4.

The main difference of this mechanics from the standard light cone gauge description consists in the fact that the gauge axis is included in the set of dynamical variables. Its position in the center-of-mass frame is described by unit norm 3-vector $\vec{e}_3$, while the gauge axis in Minkowski space is defined by light-like 4-vector $n_\mu = (N_\mu^0 - N_\mu^i e_3^i) / \sqrt{2}$, where N_μ^ν is an orthonormal tetrad introduced in Part I. In familiar terms, χ_0 generates the evolution of the string, i.e. can be used as Hamiltonian; χ_3 transforms the canonical variables of the theory in such a way that observable string's coordinates and momenta are not changed, therefore it can

¹In this paper the terms “generator of rotations”, “orbital moment” and “spin” refer to the same variable. Finally it defines the spin of elementary particles, which will result from quantization.

be interpreted as internal gauge transformation of the theory; $\chi_{1,2}$ change the orientation of the gauge axis $\vec{e}_3$ and are followed by reparametrizations of the string. Generators of Lorentz group defined in Part I produce “rigid” Lorentz transformations of the world sheet together with the gauge axis, without change of its parametrization. As a result, in quantum mechanics Lorentz group was freed of anomalies. The central problem of this approach is that the algebra of constraints χ_i in quantization acquires the same anomaly that earlier was in Lorentz group. To solve this problem, we partially eliminate the gauge degrees of freedom in the classical theory, imposing additional gauge fixing conditions to χ_i . These conditions are equivalent to fixation of gauge axis $\vec{e}_3$ with respect to other dynamical vectors in the system. As a result of this procedure, the set of constraints in the theory will be reduced, and the remainder will be anomaly-free.

Before to proceed, we will consider one remarkable particular solution in string theory.

Straight-line string is a solution, for which the string has a form of straight segment rotating about its middle at constant angular velocity in the center-of-mass frame. This solution plays a special role in string theory [2]:

- it can be quantized at arbitrary number of dimensions $d \geq 3$ without anomalies in Lorentz group;
- it belongs to a border of classical Regge-plot $P^2/2\pi \geq S$, and in quantum theory corresponds to a leading Regge-trajectory $P^2/2\pi = S$;
- it describes satisfactory the spectrum of light mesons and gives an opportunity to consider their radiative transitions.

In light cone gauge the straight-line string is represented by one-modal solution, if the gauge axis $\vec{e}_3$ is directed along the spin $\vec{S}$, orthogonally to the plane, where the straight-line string is rotating:

$$a_n = 0, \quad n \neq 1, \quad S_{\pm} = 0, \quad \frac{P^2}{2\pi} = S = S_3 = |a_1|^2. \quad (4)$$

There is a gauge-equivalent one-modal solution with excited (a_{-1}) -mode and $\vec{e}_3$ opposite to $\vec{S}$. Other directions of gauge axis for the straight-line string give infinitely-modal solutions (particular example see in Part I). Later we will use the solution (4) to study the structure of general theory in its vicinity.

Additional gauge fixing conditions we impose have a form

$$a_s + a_{-s} = 0, \quad a_s^* + a_{-s}^* = 0 \quad (5)$$

for some $s > 0$. It's easy to check that straight-line string solution (4) satisfies these conditions at $s > 1$. Later, in subsection 2.4, we will show that conditions (5) can be imposed on any solution of string theory.

The gauge fixing conditions (5) are preserved by transformations R_3 . They are not preserved by E_0 , however, there is a remainder of E_0 -symmetry, discrete transformation

$$D_{2s} : a_n \rightarrow a_n e^{-in\pi/s}$$

preserving (5): $a_s + a_{-s} \rightarrow -a_s - a_{-s}$. As a result, R_3, D_{2s} -symmetries are present in the theory after gauge fixation.

Substituting (5) to the symplectic form (2), we see that $a_{\pm s}$ -terms cancel each other: $\frac{1}{is}(da_s^* \wedge da_s - da_{-s}^* \wedge da_{-s}) =$

0. To study this property in more detail, we introduce the variables $q_1 = \text{Re}(a_s + a_{-s})/2$, $q_2 = -\text{Im}(a_s + a_{-s})/2$, $p_1 = \text{Im}(a_s - a_{-s})/2$, $p_2 = \text{Re}(a_s - a_{-s})/2$, and rewrite $a_{\pm s}$ -terms of the symplectic form as $\frac{1}{s}(dp_1 \wedge dq_1 - dp_2 \wedge dq_2)$. Gauge fixing conditions (5) are rewritten to $q_1 = q_2 = 0$, now we see that the variables $p_{1,2}$, canonically conjugated to $q_{1,2}$, drop out from the symplectic form. $p_{1,2}$ should be expressed in terms of other dynamical variables from χ -constraints, and independently on how complex these expressions could be, the symplectic structure of the mechanics remains to be simple. This exclusive property follows from the fact that two gauge fixing conditions (5) are in involution with each other: $\{a_s + a_{-s}, a_s^* + a_{-s}^*\} = 0$, i.e. generate *abelian* group of transformations.

Then we see that contribution of $a_{\pm s}$ to A_3 vanishes similarly: $\frac{1}{s}(|a_s|^2 - |a_{-s}|^2) = 0$, as a result, $a_{\pm s}$ -terms also drop out from $\chi_3 = S_3 - A_3$. This result follows from the fact that gauge fixing conditions (5) are preserved by transformation R_3 and therefore are in involution with χ_3 .

The gauge fixing conditions (5) are not in involution with χ_0 , and $a_{\pm s}$ -terms in L_0 do not vanish: $|a_s|^2 + |a_{-s}|^2 = 2|a_s|^2$. Poisson brackets of (5) with $\chi_{\pm}$ also do not vanish. We conclude that (5) are gauges for $(\chi_{\pm}, \chi_0)$ and $a_{\pm s}$ should be determined from these three constraints.

2.2 Algebraic properties of lia-lcg

Isolating contribution of $a_{\pm s}$ -oscillators in (3) and using the relation $a_{-s} = -a_s$, we have

$$a_s^2 d + \frac{1}{2} a_s a_s^* d^* + a_s f + a_s^* g + \Sigma_- - \sqrt{\frac{P^2}{2\pi}} S_- = 0, \quad (6)$$

$$a_s^{*2} d^* + \frac{1}{2} a_s a_s^* d + a_s^* f^* + a_s g^* + \Sigma_+ - \sqrt{\frac{P^2}{2\pi}} S_+ = 0,$$

$$\frac{P^2}{2\pi} = L_0^{(s)} + 2a_s a_s^*,$$

where

$$d = d_+ - d_-, \quad f = f_+ - f_-, \quad g = g_- - g_+, \quad (7)$$

$$d_+ = a_{2s}^*/s, \quad d_- = a_{-2s}^*/s, \quad L_0^{(s)} = \sum' a_k^* a_k$$

$$g_- = \sum' \frac{1}{k} a_k a_{s-k}, \quad g_+ = \sum' \frac{1}{k} a_k a_{-s-k},$$

$$f_+ = \sum' \left(\frac{1}{s} + \frac{1}{k}\right) a_k a_{s+k}^*, \quad f_- = \sum' \left(-\frac{1}{s} + \frac{1}{k}\right) a_k a_{-s+k}^*,$$

$$\Sigma_- = \sum' \frac{1}{k} a_k a_n a_{k+n}^*, \quad \Sigma_+ = \sum' \frac{1}{k} a_k^* a_n^* a_{k+n}, \quad \Sigma_-^* = \Sigma_+,$$

Here in the sums $\sum'$ terms with $a_0^{(*)}$ and $a_{\pm s}^{(*)}$ are excluded. Introducing denotations

$$\lambda = \sqrt{\frac{P^2}{2\pi}}, \quad n_s = (\lambda^2 - L_0^{(s)})/2,$$

$$k = \Sigma_- + \frac{1}{2} n_s d^* - \lambda S_-,$$

we can treat (6) as overdetermined polynomial system for (a_s, a_s^*) at given values of coefficients (d, f, g, k, n_s) :

$$a_s^2 d + a_s f + a_s^* g + k = 0, \quad (8)$$

$$a_s^{*2} d^* + a_s^* f^* + a_s g^* + k^* = 0,$$

$$a_s a_s^* - n_s = 0.$$

To solve this system we find its Groebner basis [3, 4]:

$$Gr[(8), \{a_s, a_s^*\}] = \{H, a_s^* G + F, a_s^* \tilde{G} + \tilde{F}, \dots\}, \quad (9)$$

where $H, G, F, \dots$ are polynomials of coefficients, given in Appendix 1. Groebner basis has the same set of roots as original system (8) and possesses a structure convenient for their sequential determination. The first equation $H = 0$ is a condition of consistency of the original system formulated in terms of its coefficients. The next equations give the solution of the system as rational expressions

$$a_s^* = -F/G = -\tilde{F}/\tilde{G} = \dots$$




Fig.1. Resolution of singularities in a_s^* .

Different expressions for a_s^* coincide on the surface $H = 0$, in the regions where the denominators do not vanish. In the points where the denominator in one formula vanishes, e.g. $G = 0$, we can use the other formula $a_s^* = -\tilde{F}/\tilde{G}$, provided that $\tilde{G} \neq 0$. Therefore, in such points both numerator and denominator of $a_s^* = -F/G$ must vanish simultaneously and resolution of this ambiguity gives finite value $(-\tilde{F}/\tilde{G})$. At the end of Groebner basis the polynomials of degree > 1 in a_s^* are present. Those which do not contain a_s , describe solutions for the exceptional degenerate cases when all denominators of the linear formulae vanish: $G = \tilde{G} = \dots = 0$. Those which are linear in a_s formally can be used to find a_s . The original equations are also appended at the end of Groebner basis, meaning that all solutions must be checked by substitution to the original system.

The presence of complex conjugation in (8) creates two general problems: (i) the described procedure treats (a_s, a_s^*) as independent variables $(a_s, b_s) \in \bar{C}^2$, it does not guarantee that a_s will be complex conjugated to b_s . However, the symmetry of original system with respect to the complex conjugation implies that (b_s^*, a_s^*) is also a solution of (8), possibly the other one. We are interested only by the case $(b_s^*, a_s^*) = (a_s, b_s)$. (ii) theorems [3] in the case of non-holomorphic polynomial systems do not guarantee that all solutions of $H = 0$ will be continued to full solutions of (8), obstacles are the equations of type $z^*z = -1$. To resolve both problems we substitute the expression $a_s^* = -F/G$ and its complex conjugate $a_s = -F^*/G^*$ to the original equations and explicitly check that they are satisfied on the surface $H = 0$ (after putting all terms over a common denominator we check that H is a polynomial divisor of the numerator). As a result, we obtain the solutions (a_s, a_s^*) , which are correctly transformed by complex conjugation and are defined everywhere except of zero-measure set of degenerate cases described above.

$H = 0$, the condition of consistency of the system (8), is the constraint, remaining after imposition of two gauge fixing conditions (5) to three initial constraints $\chi_0, \chi_\pm$. This condition is real-valued: $H^* = H$. Being used as Hamiltonian, it generates correct string evolution, consisting of shifts $\sigma \rightarrow \sigma + \tau$ and such reparametrizations that keep gauge fixing conditions (5) permanently satisfied. Hamiltonian is in involution with the constraint χ_3 , i.e. is preserved by transformation $R_3 : a_n \rightarrow a_n e^{-i\alpha}$, $S_- \rightarrow S_- e^{-i\alpha}$. Hamiltonian is

also preserved by transformation $D_{2s} : a_n \rightarrow a_n e^{-in\pi/s}$. Using the explicit representation of a_s from Appendix 1, one can easily verify that the properties of a_s with respect to these transformations are the same as before gauge fixation:

$$R_3(a_s) = a_s e^{-i\alpha}, \quad D_{2s}(a_s) = -a_s. \quad (10)$$

It's important to remark that these symmetries are defined by polynomial structure of the expressions: the symmetries implemented as phase rotations of oscillator variables impose certain conditions on the powers of oscillator variables in monomials. As a result, R_3, D_{2s} -symmetries can be preserved in quantum theory as well.

Generally H is 8th degree polynomial in λ , in special case $S_\pm = 0$ it degenerates to 4th degree polynomial in λ^2 . Using implicit function theorems, one can locally rewrite the constraint $H = 0$ in the form of mass-shell condition: $P^2 - \Phi = 0$, where the function Φ globally has multiple branches. Because at $s > 1$ straight-line string (4) satisfies gauge fixing conditions (5), one branch of Φ passes through the straight-line string. Further we study in details the structure of solutions in the vicinity of straight-line string.

2.3 Solutions in the vicinity of straight-line string

Lemma 1: at $s = 2$ in the vicinity of straight-line string the system (6) has unique solution, representable as C^∞ -smooth analytical function of coefficients of the system.

[Proof of the lemma can be found in Appendix 2.]

Let's fix $s = 2$. On the straight-line string solution (4) for the coefficients of the system (6) we have $f_\pm = d_\pm = \Sigma_\pm = S_\pm = 0$, in g the term $g_+ = 0$, while $g_- = a_1^2$. Let's pull off a_1 -contribution from g -terms: $g'_- = g_- - a_1^2$, $g' = g'_- - g_+$. Due to the lemma above we can represent a_2 as Taylor's expansion with respect to these coefficients. Further we will study a part of this expansion, which do not contain $S_\pm$ -terms. The case $S_\pm = 0$ possesses convenient simplifications, particularly, the first two equations in (6) define a closed system for (a_2, a_2^*) and decouple from the third one, which gives direct definition of P^2 :

$$\frac{P^2}{2\pi} = L_0^{(2)} + 2|a_2|^2.$$

Lemma 2: ($S_\pm = 0$)-terms of a_2 expansion in the vicinity of straight-line string have a form

$$a_2|_{S_\pm=0} = \sum_{n \geq 1} \frac{P_n a_1^2}{|a_1|^{4n}}, \quad (11)$$

where P_n are polynomials of a_1, Σ_-, d, f, g' and their conjugates. The polynomials possess the following properties: transformation $D_4 : a_n \rightarrow a_n i^{-n}$ preserves the monomials of P_n , transformation $R_3 : a_n \rightarrow a_n e^{-i\alpha}$ multiplies the monomials to $e^{i\alpha}$; each monomial in P_n contains, counting powers, n variables from the group $(\Sigma_-, d, f, g'$ and their conjugates) and $2(n-1)$ variables from the group (a_1, a_1^*) ; each monomial contains at least one variable $\Sigma_\pm$; numbers of variables from $(\Sigma_\pm)$ - and (d, d^*) -groups are related: $n(\Sigma) = n(d) + 1$.

Explicit expressions for the first six polynomials P_n can be found in Appendix 1. The expansion (11) has an obvious singularity at $a_1 = 0$. Later we will show that in quantum theory this singularity can be removed by a particular choice of quantum ordering.

2.4 Geometric properties of lia-lcg

In this subsection we describe in more details the transformations, generated by constraints χ_0, χ_i . At first, we introduce several definitions.

Let's consider in 3D space: smooth closed curve $\vec{Q}(\sigma)$ with marked point O and unit norm vector $\vec{e}_3$. Let's introduce variables

$$\vec{a}_n = \frac{1}{2\sqrt{2}\pi} \oint d\vec{Q}(\sigma) \cdot \exp \left[\frac{2\pi i n}{L_t} (L(\sigma) - (\vec{Q}(\sigma) - \vec{Q}(0))\vec{e}_3) \right], \quad (12)$$

where $L(\sigma)$ is a length of arc between points O and $\vec{Q}(\sigma)$ along the curve, L_t is total length of the curve. Two properties obviously follow from the definition: $\vec{a}_{-n} = \vec{a}_n^*$, $\vec{a}_0 = 0$. Let's decompose vectors $\vec{a}_n$ into the components, parallel and orthogonal to $\vec{e}_3$: $\vec{a}_n = a_{n3}\vec{e}_3 + \vec{a}_{n\perp}$, and denote real and imaginary parts of $\vec{a}_{n\perp}$ as $\vec{q}_{n\perp}$ and $\vec{p}_{n\perp}$: $\vec{a}_{n\perp} = \vec{q}_{n\perp} + i\vec{p}_{n\perp}$. Let's fix some $n = s > 0$ and write $\vec{q}_{s\perp} = \vec{q}$ and $\vec{p}_{s\perp} = \vec{p}$. Functions $\vec{q}(\vec{e}_3), \vec{p}(\vec{e}_3)$ define smooth vector fields on unit sphere of $\vec{e}_3$ (tangent to the sphere). Due to topological "hedgehog" theorem, these fields have singular points on the sphere, where corresponding field vanishes, e.g. $\vec{q} = 0$.




Fig.2. Definition of vector fields $\vec{q}(\vec{e}_3), \vec{p}(\vec{e}_3)$.

Note: the curve $\vec{Q}$ is a projection of *supporting curve*, introduced in Part I, into 3-dimensional subspace, orthogonal to total momentum P_μ . In terms of string dynamics, it is a trajectory of string's end in the center-of-mass frame. The length of this curve is equal to double mass of the string: $L_t = 2\sqrt{P^2}$. Gauge axis $\vec{e}_3$ relates the following parametrization to this curve:

$$\sigma = \frac{2\pi}{L_t} (L(\sigma) - Q_3(\sigma) + Q_3(0)), \quad (13)$$

where $Q_3 = \vec{Q}\vec{e}_3$. Now we recognize in (12) Fourier modes of function $\vec{a}(\sigma) = \vec{Q}'(\sigma)$, where σ is lcg-parameter (13). This expression is written in parametric-invariant form, as circulation integral. Vectors $\vec{a}_{n\perp}$ are related with earlier introduced oscillator variables a_n as follows:

$$\begin{aligned} \vec{a}_{n\perp} &= a_{n1}\vec{e}_1 + a_{n2}\vec{e}_2, \\ a_{n1} &= (a_n + a_{-n}^*)/2, \quad a_{n2} = i(a_n - a_{-n}^*)/2. \end{aligned}$$

Lemma 3: χ_3 generates transformations, preserving $\vec{e}_3$ and $\vec{a}_{n\perp}$; $\chi_{1,2}$ generate rotations of $\vec{e}_3$ and associated changes of $\vec{a}_{n\perp}$ according to the formula (12). Gauge fixing conditions (5) correspond to a singular point $\vec{q} = 0$. Evolution, generated by χ_0 , is represented as phase rotations $\vec{a}_{n\perp} \rightarrow \vec{a}_{n\perp} e^{-in\tau}$. During this evolution vector $\vec{\omega} = \vec{q} \times \vec{p}$ is preserved, and singular points $\vec{q} = 0$ move along zero-level curves of a function $F(\vec{e}_3) = \vec{\omega}\vec{e}_3 = 0$.

This lemma implies that gauge fixing conditions (5) can be imposed on any solution of string theory. Namely, these

conditions can be satisfied for any curve $\vec{Q}(\sigma)$, directing the light cone gauge axis $\vec{e}_3$ to a singular point $\vec{q}(\vec{e}_3) = 0$ of a vector field on the sphere, constructed in terms of this curve.

In general position the vector fields on the sphere have even number of singular points, which is ≥ 2 . Non-degenerate singular points of 2D vector fields are [5]




Fig.3. Singular points of 2D vector field.

For each type *index* of singularity is defined as algebraic number of rotations of vector $\vec{q}(\vec{e}_3)$, when $\vec{e}_3$ passes around singular point (> 0 , if directions of rotations of $\vec{q}$ and $\vec{e}_3$ coincide; < 0 otherwise). For (center, node, focus) it is $+1$, for saddle -1 . The sum of all indices is equal to Euler characteristic of the surface $V - E + F$: (num. of vertices) - (num. of edges) + (num. of faces) for any tessellation of the surface into faces, which for the sphere equals 2. Thus, generic vector field on a sphere has 2 singular points of index $+1$ and arbitrary number of self-compensating pairs $(+1, -1)$.

Presence of several singular points $\vec{q}(\vec{e}_3) = 0$ leads to the fact that an orbit of a gauge group generated by χ -constraints intersects a surface of gauge fixing conditions (5) in several points of the phase space. This phenomenon is also encountered in the theory of non-abelian gauge fields, where it has been studied by V.N.Gribov [6]. We will call such points *Gribov's copies*.




Fig.4. Gribov's copies: the orbit of a gauge group $o(G)$ intersects the surface of gauge condition $\xi = 0$ in several points.

In our case the copies comprise different sets of oscillator variables $\{a_n\}$, reproducing the same curve $\vec{Q}(\sigma)$ (differently parameterized), i.e. physically correspond to the same string's motion. This property indicates remaining *discrete* gauge equivalence in the theory. To identify the equivalent states, in classical theory the phase space should be factorized with respect to this symmetry. Analogous procedures can be applied in quantum mechanics, e.g. by constructing irreducible representations for operators possessing this discrete symmetry and formulating respective selection rules.

During the evolution Gribov's copies move along closed loops, constructed by lifting of zero-level curves $F(\vec{e}_3) = 0$ to the space of oscillator variables. In more details, this construction is performed as follows: vector $\vec{e}_3(\tau)$ moves along the contour $F(\vec{e}_3) = 0$ on the sphere, defined for a given value s , while for $n \neq s$ expression (12) defines time-dependent vector fields $\vec{a}_{n\perp}(\vec{e}_3, \tau)$, where we substitute $\vec{e}_3 = \vec{e}_3(\tau)$. Note that corresponding vector fields $\vec{q}_{n\perp}, \vec{p}_{n\perp}$ generally do not vanish on these loops. The set of Gribov's copies possesses D_{2s} -symmetry, however in general case this symmetry does not exhaust the discrete gauge equivalence.

Lemma 4: during the period of evolution $\Delta\tau = 2\pi$ singular points $\vec{q} = 0$ return to the initial position $2s$ times. The lifting of the contour $F(\vec{e}_3) = 0$ to the space of oscillator

variables has at least $2s$ Gribov's copies, related by D_{2s} -symmetry.

Below we study the structure of Gribov's copies in the vicinity of straight-line string solution (4) and its $\vec{e}_3 \rightarrow -\vec{e}_3$ companion. For these purpose we investigate the expansion of $\vec{q}(\vec{e}_3)$ for this solution in the vicinity of northern and southern poles of the sphere $\vec{e}_3 = \pm \vec{S}/|\vec{S}|$.

Lemma 5: $\vec{q}(\vec{e}_3)$ for straight-line string solution at $s = 1$ has no singularities in the vicinity of the poles; at $s > 1$ has in each pole a singularity, which does not move during the evolution, and at $s = 2$ is a saddle point, at $s > 2$ has degenerate type.

Lemma 6: Solutions, close to (4), have at $s = 2$ a saddle point moving along a small loop in the vicinity of the northern pole. During the period of evolution $\Delta\tau = 2\pi$ the saddle point performs 4 turns. Associated lifting has exactly 4 Gribov's copies, related by D_4 -symmetry.

Analogous structure of Gribov's copies is established for the companion solution near the southern pole. Solutions close to straight-line string at $s = 2$ should also have other singularities faraway of the poles (saddle points have index -2 , while the total index of singular points on the sphere should be $+2$). Further in quantum mechanics we will use the expansion (11) in the vicinity of solution (4), corresponding to the northern pole singularity.

Summary of classical mechanics: the phase space $(Z, P, a_n, a_n^*, \vec{S}, \vec{e}_i)$ possesses trivial topology. Hamiltonian H is globally defined in it as a polynomial function (Appendix 1). The following two features can be considered as topological defects: (i) the expression of mass squared $P^2 = \Phi$ from the constraint $H = 0$ is multi-valued; (ii) there is a discrete remainder of original gauge invariance, which includes D_{2s} -symmetry, but generally is not exhausted by it. In the vicinity of straight-line string solution (4) at $s = 2$ the branch of $P^2 = \Phi$ is regular, and the discrete gauge invariance is exhausted by D_4 -symmetry.

3 Quantum mechanics

3.1 Canonical operators

$$\begin{aligned} [Z_\mu, P_\nu] &= -ig_{\mu\nu}, \\ [a_k, a_n^+] &= k\delta_{kn}, \quad k, n \neq 0, \pm s, \\ [S^i, S^j] &= i\epsilon_{ijk}S^k, \quad [S^i, e_j^k] = i\epsilon_{ikl}e_j^l, \\ [S_i, S_j] &= -i\epsilon_{ijk}S_k, \quad [S_i, e_j^k] = -i\epsilon_{ijl}e_l^k, \\ [S^i, S_j] &= 0, \quad e_i^k e_j^k = \delta_{ij}, \quad S_i = e_i^j S^j. \end{aligned} \quad (14)$$

The space of states is a direct product of three components:

Space of functions $\Psi(P)$ with the definition $Z = -i\partial/\partial P$.

Fock space with a vacuum

$$a_k|0\rangle = 0, \quad k > 0, \quad a_k^+|0\rangle = 0, \quad k < 0 \quad (15)$$

and states created from vacuum by operators

$$\begin{aligned} |\{n_k\}\rangle &= \prod_{k>0, k \neq s} \frac{1}{\sqrt{k^{n_k} n_k!}} (a_k^+)^{n_k} \cdot \\ &\cdot \prod_{k<0, k \neq -s} \frac{1}{\sqrt{(-k)^{n_k} n_k!}} (a_k)^{n_k} |0\rangle. \end{aligned}$$

For instance, we will write $|1_1 2_{-1}\rangle = \frac{1}{\sqrt{2}} a_1^+ a_{-1}^2 |0\rangle$ etc. So defined state vectors have positive norm. Occupation numbers

$$n_k = \frac{1}{|k|} : a_k^+ a_k : = \frac{1}{|k|} \cdot \begin{cases} a_k^+ a_k, & k > 0 \\ a_k a_k^+, & k < 0 \end{cases} = 0, 1, 2, \dots$$

Note: one-component oscillator variables a_k used throughout this paper are related with commonly applied two-component oscillators $a_{k1,2}$ by the formulae of subsection 2.4, which on quantum level become: $a_k = a_{k1} - ia_{k2}$, $a_k^+ = a_{-k1} + ia_{-k2}$. The inverse formulae are $a_{k1} = (a_k + a_{-k}^+)/2$, $a_{k2} = i(a_k - a_{-k}^+)/2$. Here taking *two sets* of oscillator variables a_{ki} , $i = 1, 2$ with $a_{ki}^+ = a_{-ki}$ we construct *one set* without this property. Usage of one-component oscillators simplifies the algebra. It's easy to verify that definition of vacuum (15) is equivalent to a standard one $a_{ki}|0\rangle = 0, k > 0$ and the states $|\{n_k\}\rangle$ are the linear combinations of $\prod_{k>0} (a_{k1}^+)^{n_{k1}} (a_{k2}^+)^{n_{k2}} |0\rangle$.

Quantum top: the space of states is formed by functions $\Psi(e)$, $e_i^j \in SO(3)$. Because the rotation group is not simply connected, $\pi_1(SO(3)) = \mathbf{Z}_2$, two representations are possible: single- and double-valued [7]. Spin is defined as differential operator $S^i = -i\epsilon_{ijk}e_i^j \partial/\partial e_l^k$, while the projection of spin onto the coordinate system $\vec{e}_i$ is $S_i = i\epsilon_{ijk}e_j^l \partial/\partial e_l^k$. Operator S^i generates the rotation of the coordinate system $\vec{e}_i$ in external space, while S_i generates the rotations about the axes $\vec{e}_i$. These transformations act on different indices in e_i^k . They commute, therefore S^3 and S_3 are simultaneously observable.

The basis in the space of states is formed by Wigner's $\mathcal{D}$ -functions

$$|SS_3S^3\rangle = \mathcal{D}_{S_3S^3}^S(e), \quad S_3, S^3 = -S, -S+1 \dots S,$$

where S are integer for single-valued representation and half-integer for double-valued one. Further a constraint $S_3 = A_3 \in \mathbf{Z}$ will select only integer S -values².

Operator $S^2 = S^i S^i = S_i S_i$ has eigenvalues $S(S+1)$. The constraints, being functions of S_i and oscillator variables, commute with S^2 , as a result, the determination of mass spectrum can be performed separately for each S value. Particularly, spin singlet $S = 0$ corresponds to state vectors independent on e_i^j . On such states all $S^i = S_i = 0$.

Constraints $\chi_3|\Psi\rangle = H|\Psi\rangle = 0$ are imposed on the states in the spirit of Dirac's quantization of constrained theories. Here $\chi_3 = S_3 - A_3^{(s)}$,

²However, using formal modifications $A_3 \rightarrow A_3 + 1/2$, classically vanishing after the recovery of Planck's constant, it's possible to obtain half-integer spin as well. The question on possibility of inclusion of half-integer S -values in this mechanics should be better investigated.

$$A_3^{(s)} = \sum_{k \neq 0, \pm s} \frac{1}{k} : a_k^+ a_k : = \sum_{k \neq 0, \pm s} \text{sign } k \cdot n_k,$$

$$L_0^{(s)} = \sum_{k \neq 0, \pm s} : a_k^+ a_k : = \sum_{k \neq 0, \pm s} |k| \cdot n_k.$$

In the next subsection we will construct two versions of Hamiltonian H : polynomial of $\lambda = \sqrt{P^2/2\pi}$ and $P^2 - \Phi$ with Φ is defined by Taylor's series (11). For the operators entering in H : $d_\pm, f_\pm, g_\pm, \Sigma_\pm$ and their conjugates we reserve a term *elementary operators*. They can be defined by their classical expressions (7), with only replacement of complex conjugation to Hermitian one. These definitions have no ordering ambiguities. For the states with the finite number of filled modes the matrix elements of elementary operators are described by finite sums.

Symmetries R_3, E_0, D_{2s} in quantum theory are generated by operators

$$R_3 = e^{i\chi_3 \alpha}, \quad E_0 = e^{iL_0^{(s)} \tau}, \quad D_{2s} = e^{iL_0^{(s)} \pi/s}.$$

It's convenient to introduce the notion of $(\Delta A_3, \Delta L_0)$ -charges for operators satisfying the relations:

$$[A_3^{(s)}, op] = \Delta A_3 \cdot op, \quad [L_0^{(s)}, op] = \Delta L_0 \cdot op,$$

i.e. the operators, which increase or decrease the quantum numbers $(A_3^{(s)}, L_0^{(s)})$ by $(\Delta A_3, \Delta L_0)$ units:

$$op |A_3^{(s)}, L_0^{(s)}\rangle = |A_3^{(s)} + \Delta A_3, L_0^{(s)} + \Delta L_0\rangle.$$

Particularly, $\Delta A_3(a_n) = -1$, $\Delta L_0(a_n) = -n$. Hermitian conjugation of operators reverses their charges. The elementary operators have definite $(\Delta A_3, \Delta L_0)$ -charges, see Table 1. The discrete symmetry D_{2s} is also characterized by the charge $Q = L_0^{(s)} \bmod 2s$, so that the subspaces with given Q are eigenspaces $D_{2s} |Q\rangle = e^{i\pi Q/s} |Q\rangle$, and D_{2s} -symmetric operators keep these subspaces invariant.

Because the symmetries of classical theory in our case are defined by its polynomial structure, they are preserved on the quantum level. Particularly, quantum Hamiltonian possesses R_3, D_{2s} -symmetries, as a result, its non-zero matrix elements form blocks, located on $A_3^{(s)}$ Q -diagonal: $\langle A_3^{(s)} Q | H | A_3^{(s)} Q \rangle$.

3.2 Quantum Hamiltonian

Hamiltonian H can be defined directly by the expressions of Appendix 1, where we only need to fix a proper ordering (Table 1). As a result, we obtain a polynomial spectral problem [8] of the form $H(\lambda)\Psi_\lambda = 0$, where $H(\lambda) = \sum h_n \lambda^n$, the coefficients h_n are Hermitian operators and $\lambda = \sqrt{P^2/2\pi}$ is a spectral parameter. The problems of these kind are equivalent to a standard linear spectral problem in larger space: $(M - \lambda)\tilde{\Psi}_\lambda = 0$, where

$$M = \begin{pmatrix} 0 & 1 & 0 & \dots & 0 \\ 0 & 0 & 1 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & 0 & \dots & 1 \\ -\tilde{h}_0 & -\tilde{h}_1 & -\tilde{h}_2 & \dots & -\tilde{h}_{l-1} \end{pmatrix},$$

l is a degree of polynomial $H(\lambda)$ and $\tilde{h}_n = h_l^{-1} h_n$, providing that h_l^{-1} exists. Using the regularizations based on the restriction of operators to finite-dimensional spaces $|L_0^{(s)} \leq N\rangle$ (described below), the problem is reduced to a spectral analysis of a large matrix, which can be performed with the aid of standard computational algorithms [9]. Due to the block structure of Hamiltonian, λ spectrum can be defined separately in each $A_3^{(s)}$ Q -block.

The following circumstance is rarely remarked. Generally Hamiltonian H can depend on P^2 non-linearly. Classically one can resolve the constraint $H = 0$ with respect to P^2 , and obtain a locally equivalent theory. On the quantum level these theories have principal differences: the first one creates non-linear spectral problem, while the second one gives the spectral problem $(P^2 - \Phi)\Psi = 0$, linear in terms of P^2 . The last problem has relatively simple structure of solutions: Hermitian operator Φ possesses complete orthonormal basis of eigenvectors and real eigenvalues, representing the mass spectrum. In the polynomial problem Hermitian operator $H(\lambda)$ at fixed λ also has real eigenvalues and complete orthonormal basis of eigenvectors: $H(\lambda)\Psi_{\lambda,\mu} = \mu(\lambda)\Psi_{\lambda,\mu}$. Here the eigenvalues and eigenvectors are the functions of λ . However, the vectors satisfying $H(\lambda)\Psi_\lambda = 0$, defined by the roots of the equation $\mu(\lambda) = 0$, should not necessarily form a complete basis. For finite-dimensional matrix systems the number of real roots of this equation can be less than the dimension of the space, it also can be greater than the dimension. Furthermore, the vectors $\Psi_\lambda = 0$ for different λ generally are not orthogonal.

The last property can be improved. In a particular case when Ψ_λ for the real roots λ form a complete basis, we can construct the matrix $Y = X\Lambda X^{-1}$, where the columns of X are vectors Ψ_λ and $\Lambda = \text{diag } \lambda$. It's easy to verify that under these conditions Y is a solution of matrix equation $\sum h_n Y^n = 0$, where Y^n stand on the right of h_n , while Y^+ is a similar solution from the left: $\sum Y^{+n} h_n = 0$. In the classical limit both Y and Y^+ correspond to a real root λ of classical equation $\sum h_n \lambda^n = 0$. Operator $(Y + Y^+)/2$ is Hermitian quantum analog of λ . It has orthogonal system of eigenvectors and can be used to define the mass spectrum.

The problem with the number of eigenvalues is more serious. If the classical equation $H(\lambda) = 0$ has several real roots λ_i , one can represent each root by a separate function on the phase space. In quantum mechanics each root becomes Hermitian operator possessing a complete system of eigenvectors. However, in the solutions of polynomial spectral problem the corresponding eigensystems are mixed together and their separation appears to be non-trivial problem. Theory of matrix polynomials factorization [8] possesses sufficient potential to solve this problem. Particularly, it introduces sign characteristics of eigenvalues, which under certain conditions allow to separate them completely. On the other hand, the resolution of constraint $H = 0$ to $P^2 = \Phi$ and

Table 1: $L_0^{(2)}$ -normally ordered elementary operators

	a_1^+	$(d_-)^+$	d_+	$(f_-)^+$	f_+	$(g_-')^+$	g_+	Σ_-	Σ_+	$(g_+)^+$	g_-'	$(f_+)^+$	f_-	$(d_+)^+$	d_-	a_1
$\Delta L_0^{(2)}$	1	4	4	2	2	2	2	0	0	-2	-2	-2	-2	-4	-4	-1
$\Delta A_3^{(2)}$	1	-1	1	0	0	2	-2	-1	1	2	-2	0	0	-1	1	-1

separation of Φ branches on the classical level provides an alternative approach to this problem. Particularly, the spectrum of P^2 in the vicinity of straight-line string can be studied using the expansion (11). In this paper we will consider this problem for a particular case of spin singlet $S = 0$.

3.3 Solutions in the vicinity of straight-line string

Let's fix $s = 2$ and put $S_3 = S_{\pm} = 0$ in the formulae above. Quantum analog of (11) is constructed as follows:

$$a_2 = \sum_{n \geq 1} : P_n a_1^2 : D^{-2n}, \quad (16)$$

where $D = a_1^+ a_1 + 1/2 = (a_1^+ a_1 + a_1 a_1^+)/2$.

Polynomials P_n are defined by expressions of Appendix 1, where we substitute the definitions (7) of elementary operators and fix the ordering, shown in Table 1. This ordering puts $L_0^{(2)}$ -lowering elementary operators to the right from $L_0^{(2)}$ -raising ones, thus providing better convergence properties for the expansion (16). Particularly, the matrix elements of a_2 between the states with finite $L_0^{(2)}$ are given by finite sums, and large n terms of (16) contribute only to the matrix elements with large $L_0^{(2)}$, due to the following lemma.

Lemma 7:

$\langle L_0^{(2)} = N_1 | : P_n : | L_0^{(2)} = N_2 \rangle = 0$, if $4(n-1) > N_1 + N_2$.

Square-mass operator is given by

$$\frac{P^2}{2\pi} = L_0^{(2)} + 2a_2^+ a_2. \quad (17)$$

One can also include here classically vanishing constant term: $P^2 \rightarrow P^2 + c$. In this definition (at $c > 0$) the operator P^2 is Hermitian and positively defined³.

The introduced ordering possesses the other feature convenient for computations: the matrix elements of the normally ordered operators restricted to finite-dimensional subspaces, behave regularly when the dimension of subspaces increases. Namely, the matrix in larger space includes the matrix in smaller space as an exact submatrix. Without normal ordering this simple behavior would be violated.

Lemma 8: let $op_i(N) = \langle L_0^{(2)} \leq N | op_i | L_0^{(2)} \leq N \rangle$ be restrictions of elementary operators op_i to finite-dimensional space $L_0^{(2)} \leq N$. Let $: P(op_i) :$ be $L_0^{(2)}$ -normally ordered polynomial of op_i . Then (i) at $N_1 < N_2$ the matrix $: P(op_i(N_1)) :$ is a submatrix of $: P(op_i(N_2)) :$ and (ii) $: P(op_i(N)) := \langle L_0^{(2)} \leq N | : P(op_i) : | L_0^{(2)} \leq N \rangle$.

Practically, we compute the matrix elements of elementary operators in $L_0^{(2)} \leq N$ subspace. The dimension of this subspace is rapidly increasing with N , e.g. $\dim = 106587$

³Note that other possible definition $P^2/2\pi = L_0^{(2)} + 2 : a_2^+ a_2 :$ does not have this property.

for $N = 25$, so that the elementary operators are represented as matrices of very large size ($\dim \times \dim$), see Appendix 3. The matrices have noticeable block structure, corresponding to their $(A_3^{(2)}, L_0^{(2)})$ -charge properties. As a result, non-zero matrix elements of a_2 , necessary for computation of P^2 at $S = 0$, are located in the blocks $\langle A_3 = -1, Q + 2 | a_2 | A_3 = 0, Q \rangle$, while P^2 itself is located in $\langle A_3 = 0, Q | P^2 | A_3 = 0, Q \rangle$. In addition to these properties we use the fact that elementary matrices inside $(A_3^{(2)}, L_0^{(2)})$ -blocks are very sparse (at large N their non-zero content occupies less than 1% of the blocks), and implement special algorithms for sparse block matrix computations, described in more details in Appendix 3. Finally, we determine the spectrum of $P^2/2\pi$ up to the value $N = 28$ ($\dim = 260256$) and the number of terms in expansion (16) up to $n = 6$. The initial part of P^2 spectrum is rapidly stabilized when n is increasing. Particularly, the region $P^2/2\pi \leq 10$ is defined by $n \leq 3$ terms, while $n > 3$ corrections influence the higher P^2 values only. Figure 5 shows the resulting spectrum of P^2 , for separate Q -values and superimposed.



 Fig.5. Spectrum ($P^2/2\pi, S = 0$).

 Table 2: eigenvectors ($P^2/2\pi \leq 5, S = 0$).

$P^2/2\pi = L_0^{(2)}$	$\{ n_k\rangle\}$
0	$ 0\rangle$
2	$ 1_1 1_{-1}\rangle$
4	$ 2_1 2_{-1}\rangle, 1_3 1_{-1}\rangle, 1_1 1_{-3}\rangle$
5	$ 1_4 1_{-1}\rangle, 1_1 1_{-4}\rangle$

The spectrum contains degenerate eigenvalues, whose multiplicities are shown on the figure by numbers in brackets (non-degenerate eigenvalues have no these numbers). Black points display the states annihilated by operator a_2 , which are simultaneously the eigenvectors of $L_0^{(2)}$ and therefore have integer $P^2/2\pi$. The beginning of the spectrum consists only of such states, as the following lemma shows.

Lemma 9: a_2 annihilates $|L_0^{(2)} \leq 5\rangle$.

At $P^2/2\pi \geq 6$ the eigenvalues of $L_0^{(2)}$ start to split. Particularly, one state with $L_0^{(2)} = 6$ goes to $P^2/2\pi = 8.16792$, two states with $L_0^{(2)} = 8$ become $P^2/2\pi = 9.0107, 10.018$ etc. Such states (displayed by white points on fig.5) correspond to the excited a_2 -modes. The values of $P^2/2\pi$ for

these states are not integer (however, some of them are numerically close to integer). Many eigenvectors of $L_0^{(2)} \geq 6$ are also annihilated by a_2 , their $P^2/2\pi$ are degenerate and integer. This degeneracy can be removed by further model corrections, such as spin-orbital interaction discussed in [2].

Subspaces with different Q in quantum theory play a role of factor-spaces with respect to D_4 -symmetry, which is satisfied exactly for $Q = 0$: $D_4\Psi = \Psi$, and in projective sense for $Q \neq 0$: $D_4\Psi \sim \Psi$. Quantum Hamiltonians of interaction, possessing D_4 -symmetry, keep these subspaces invariant. Hamiltonians, violating this symmetry, create transitions between different Q -sectors. If the violation of D_4 -symmetry is small, the particles on the first daughter Regge trajectories will be quasi-stable.

Summary of quantum mechanics: representation of canonical operators (14) is straightforward. Quantum Hamiltonian can be defined either as polynomial function (Appendix 1) or as expansion (16) in the vicinity of straight-line string, both properly ordered. In the first definition a non-linear spectral problem appears, with associated separation task for the roots of characteristic equation. The second definition leads to the ordinary linear spectral problem, which allows, particularly, to determine the mass spectrum for the states of spin singlet $S = 0$.

A paper devoted to systematic study of Gribov's copies in this mechanics and determination of mass spectrum outside of the spin singlet will be published.

References

- [1] I. Nikitin, String theory in Lorentz-invariant light cone gauge, hep-th/9906003
- [2] E.B. Berdnikov, G.G. Nanobashvili, G.P. Pron'ko, The Relativistic Theory for Principal Trajectories and Electromagnetic Transitions of Light Mesons, Int. J. Mod. Phys. A 1993. V.8. N14. P.2447; V.8. N15. P.2551.
- [3] D. Cox, J. Little, D. O'Shea, Ideals, Varieties, and Algorithms, Springer 1998.
- [4] S. Wolfram, The Mathematica^R Book, Cambridge University Press 1999.
- [5] B.A. Dubrovin, S.P. Novikov, A.T. Fomenko, Modern Geometry, Moscow, Nauka 1979.
- [6] V.N. Gribov, Quantization of non-abelian gauge theories, Nucl.Phys. B139. 1978. P.1.
- [7] H. Weyl, The theory of groups and quantum mechanics, New York: Dover, 1950.
- [8] I. Gohberg, P. Lancaster, L. Rodman, Matrix Polynomials, Academic Press, 1982.
- [9] J.H. Wilkinson, C. Reinsch, Linear Algebra, vol.2 of Handbook for Automatic Computations, New York: Springer Verlag 1971.
- [10] V.I. Arnold, Lectures on bifurcations and versal families, Rus.Math.Surv. (Uspekhi Mat. Nauk) 1972. V.27 P.119.
- [11] W.H. Press, S.A. Teukolsky, W.T. Vetterling, B.P. Flannery, Numerical Recipes in C, Cambridge University Press, 1988.
- [12] R. Davies, NewMat C++ Matrix Class, <http://ideas.uqam.ca/ideas/data/Softwares/codccplusnewmat.html>

Appendix 1: the polynomials, entering to the definition of Groebner's basis (9) of the system (8):

$$H = -f^* g g^{*2} k^2 - f g^2 g^* k^{*2} - d f^* g k k^{*2} - d^* f g^* k^2 k^* + d^* g^{*2} k^3 + d g^2 k^{*3} + d^* f^2 g^* k n_s + d f f^{*2} g k^* n_s + 2 d^* f^2 g g^* k^* n_s + 2 d f^{*2} g g^* k n_s - 3 d^* f g g^{*2} k n_s - 3 d f^* g^2 g^* k^* n_s - 2 d d^* f^* g^* k^2 n_s - 2 d d^* f g k^{*2} n_s - d^2 f^{*3} g n_s^2 - d^{*2} f^3 g^* n_s^2 + d^2 d^* f^{*2} k n_s^2 + d d^{*2} f^2 k^* n_s^2 + 3 d d^{*2} f g^* k n_s^2 + 3 d^2 d^* f^* g k^* n_s^2 + f f^* g g^* k k^* + g^2 g^{*2} k k^* + d d^* k^2 k^{*2} - g^3 g^{*3} n_s + 2 f f^* g^2 g^{*2} n_s - f^2 f^{*2} g g^* n_s - d d^* f f^* k k^* n_s + d d^* g g^* k k^* n_s - d d^* f f^* g g^* n_s^2 + 3 d d^* g^2 g^{*2} n_s^2 - 2 d^2 d^{*2} k k^* n_s^2 - 3 d^2 d^{*2} g g^* n_s^3 - d^2 d^{*2} f f^* n_s^3 + d^3 d^{*3} n_s^4$$

$$F = -f^{*3} g^2 g^* k n_s + 2 d^* f^{*2} g g^* k^2 n_s - d^{*2} f^* g^* k^3 n_s + f^{*2} g^3 g^* k^* n_s - d^{*2} g g^* k^2 k^* n_s - d^* g^3 g^* k^{*2} n_s + d f^{*4} g^2 n_s^2 - d^* f^* g^3 g^{*2} n_s^2 - 2 d d^* f^{*3} g k n_s^2 - 2 d^{*2} f f^* g g^* k n_s^2 + d^{*2} g^2 g^{*2} k n_s^2 + d d^{*2} f^{*2} k^2 n_s^2 + 2 d^{*3} f g^* k^2 n_s^2 - 3 d d^* f^{*2} g^2 k^* n_s^2 + 2 d^{*2} f g^2 g^* k^* n_s^2 + 4 d d^{*2} f^* g k k^* n_s^2 - d d^{*3} k^2 k^* n_s^2 + d d^{*2} g^2 k^{*2} n_s^2 + 2 d d^{*2} f f^{*2} g n_s^3 - d^{*3} f^2 g g^* n_s^3 + 2 d d^{*2} f^* g^2 g^* n_s^3 - 2 d d^{*3} f f^* k n_s^3 - 2 d d^{*3} g g^* k n_s^3 - 2 d d^{*3} f g k^* n_s^3 + d d^{*4} f^2 n_s^4 - d^2 d^{*3} f^* g n_s^4 + d^2 d^{*4} k n_s^4$$

$$G = f^{*3} g^2 k k^* - 2 d^* f^{*2} g k^2 k^* + d^{*2} f^* k^3 k^* - f^{*2} g^3 k^{*2} + d^{*2} g k^2 k^{*2} + d^* g^3 k^{*3} - f f^{*4} g^2 n_s + f^{*3} g^3 g^* n_s + 2 d^* f f^{*3} g k n_s - 3 d^* f^{*2} g^2 g^* k n_s - d^{*2} f f^{*2} k^2 n_s + 3 d^{*2} f^* g g^* k^2 n_s - d^{*3} g^* k^3 n_s + 3 d^* f f^{*2} g^2 k^* n_s - 2 d^{*2} f f^* g k k^* n_s - d^{*3} f k^2 k^* n_s - 3 d^{*2} f g^2 k^{*2} n_s - 2 d^{*2} f^2 f^{*2} g n_s^2 + 2 d^{*3} f^2 f^* k n_s^2 + 3 d^{*3} f^2 g k^* n_s^2 - d^{*4} f^3 n_s^3$$

The polynomials, entering to the expansion (11) of variable $a_2|_{S_{\pm}=0}$ in the vicinity of straight-line string solution (4):

$$P_1 = -\Sigma_+,$$

$$P_2 = a_1^{*2} f^* \Sigma_- + a_1^2 g'^* \Sigma_+,$$

$$2P_3 = -2 a_1^{*4} f^* g' \Sigma_- - 2 a_1^2 a_1^{*2} f^* g'^* \Sigma_- - 2 a_1^{*4} d^* \Sigma_-^2 - 2 a_1^2 a_1^{*2} f f^* \Sigma_+ - 2 a_1^4 g'^* \Sigma_+ - a_1^2 a_1^{*2} d \Sigma_- \Sigma_+,$$

$$2P_4 = 2 a_1^2 a_1^{*4} f f^{*2} \Sigma_- + 2 a_1^{*6} f^* g'^2 \Sigma_- + 2 a_1^2 a_1^{*4} f^* g' g'^* \Sigma_- + 2 a_1^4 a_1^{*2} f^* g'^* \Sigma_- + a_1^2 a_1^{*4} d f^* \Sigma_-^2 + 4 a_1^{*6} d^* g' \Sigma_-^2 + 2 a_1^2 a_1^{*4} d^* g'^* \Sigma_-^2 + 2 a_1^2 a_1^{*4} f f^* g' \Sigma_+ + 4 a_1^4 a_1^{*2} f f^* g'^* \Sigma_+ + 2 a_1^6 g'^* \Sigma_+ + 4 a_1^2 a_1^{*4} d^* f \Sigma_- \Sigma_+ + a_1^2 a_1^{*4} d^* f^* \Sigma_- \Sigma_+ + a_1^2 a_1^{*4} d g' \Sigma_- \Sigma_+ + 2 a_1^4 a_1^{*2} d g'^* \Sigma_- \Sigma_+ + a_1^4 a_1^{*2} d f \Sigma_+^2 + 2 a_1^4 a_1^{*2} d f^* \Sigma_+^2,$$

$$4P_5 = -8 a_1^2 a_1^{*6} f f^{*2} g' \Sigma_- - 4 a_1^{*8} f^* g'^3 \Sigma_- - 8 a_1^4 a_1^{*4} f f^{*2} g'^* \Sigma_- - 4 a_1^2 a_1^{*6} f^* g'^2 g'^* \Sigma_- - 4 a_1^4 a_1^{*4} f^* g' g'^* \Sigma_- - 4 a_1^6 a_1^{*2} f^* g'^* \Sigma_- - 12 a_1^2 a_1^{*6} d^* f f^* \Sigma_-^2 - 2 a_1^2 a_1^{*6} d^* f^{*2} \Sigma_-^2 - 4 a_1^2 a_1^{*6} d f^* g' \Sigma_-^2 - 12 a_1^{*8} d^* g'^2 \Sigma_-^2 - 4 a_1^4 a_1^{*4} d f^* g'^* \Sigma_-^2 - 8 a_1^2 a_1^{*6} d^* g' g'^* \Sigma_-^2 - 4 a_1^4 a_1^{*4} d^* g'^* \Sigma_-^2 - 2 a_1^2 a_1^{*6} d d^* \Sigma_-^2 - 2 a_1^2 a_1^{*6} d f^* \Sigma_-^3 - 4 a_1^4 a_1^{*4} f^2 f^{*2} \Sigma_+ - 4 a_1^2 a_1^{*6} f f^* g'^2 \Sigma_+ - 8 a_1^4 a_1^{*4} f f^* g' g'^* \Sigma_+ - 12 a_1^6 a_1^{*2} f f^* g'^* \Sigma_+ - 4 a_1^8 g'^* \Sigma_+ - 8 a_1^4 a_1^{*4} d f f^* \Sigma_- \Sigma_+ - 8 a_1^4 a_1^{*4} d f^* \Sigma_- \Sigma_+ - 8 a_1^4 a_1^{*4} d f^{*2} \Sigma_- \Sigma_+ - 16 a_1^2 a_1^{*6} d^* f g' \Sigma_- \Sigma_+ - 4 a_1^2 a_1^{*6} d^* f^* g'^* \Sigma_- \Sigma_+ - 4 a_1^4 a_1^{*4} d g' g'^* \Sigma_- \Sigma_+ - 6 a_1^6 a_1^{*2} d g'^* \Sigma_- \Sigma_+ - a_1^4 a_1^{*4} d^2 \Sigma_-^2 \Sigma_+ - 4 a_1^2 a_1^{*6} d^{*2} \Sigma_-^2 \Sigma_+ - 4 a_1^4 a_1^{*4} d^* f^2 \Sigma_+^2 - 2 a_1^4 a_1^{*4} d^* f f^* \Sigma_+^2 - 2 a_1^4 a_1^{*4} d f g' \Sigma_+^2 - 4 a_1^4 a_1^{*4} d f^* g' \Sigma_+^2 - 6 a_1^6 a_1^{*2} d f g'^* \Sigma_+^2 - 12 a_1^6 a_1^{*2} d f^* g'^* \Sigma_+^2 - 9 a_1^4 a_1^{*4} d d^* \Sigma_- \Sigma_+^2 - 2 a_1^6 a_1^{*2} d^2 \Sigma_+^3,$$

$$4P_6 = 4 a_1^4 a_1^{*6} f^2 f^{*3} \Sigma_- + 12 a_1^2 a_1^{*8} f f^{*2} g'^2 \Sigma_- + 4 a_1^{*10} f^* g'^4 \Sigma_- + 16 a_1^4 a_1^{*6} f f^{*2} g' g'^* \Sigma_- + 4 a_1^2 a_1^{*8} f^* g'^3 g'^* \Sigma_- + 12 a_1^6 a_1^{*2} f f^{*2} g'^* \Sigma_- + 4 a_1^4 a_1^{*4} f^* g'^2 g'^* \Sigma_- + 4 a_1^6 a_1^{*2} f^* g' g'^* \Sigma_- + 4 a_1^8 a_1^2 f^* g'^* \Sigma_- + 6 a_1^4 a_1^{*4} d f f^{*2} \Sigma_-^2 + 4 a_1^4 a_1^{*6} d f^{*3} \Sigma_-^2 + 36 a_1^2 a_1^{*8} d^* f f^* g' \Sigma_-^2 + 6 a_1^2 a_1^{*8} d^* f^{*2} g' \Sigma_-^2 + 6 a_1^2 a_1^{*8} d f^* g'^2 \Sigma_-^2 + 16 a_1^{*10} d^* g'^3 \Sigma_-^2 + 24 a_1^4 a_1^{*6} d^* f f^* g'^* \Sigma_-^2 + 4 a_1^4 a_1^{*6} d^* f^{*2} g'^* \Sigma_-^2 + 8 a_1^4 a_1^{*6} d f^* g' g'^* \Sigma_-^2 + 12 a_1^2 a_1^{*8} d^* g'^2 g'^* \Sigma_-^2 + 6 a_1^6 a_1^{*2} d f^* g'^* \Sigma_-^2 + 8 a_1^4 a_1^{*4} d^* g' g'^* \Sigma_-^2 + 4 a_1^6 a_1^{*2} d^* g'^* \Sigma_-^2 + 8 a_1^2 a_1^{*8} d^{*2} f \Sigma_-^3 + a_1^4 a_1^{*6} d^2 f \Sigma_-^3 + 6 a_1^2 a_1^{*8} d^{*2} f^* \Sigma_-^3 + 6 a_1^2 a_1^{*8} d^* d^* g' \Sigma_-^3 + 4 a_1^4 a_1^{*6} d d^* g'^* \Sigma_-^3 + 8 a_1^4 a_1^{*6} f^2 f^{*2} g' \Sigma_+ + 4 a_1^2 a_1^{*8} f f^* g'^3 \Sigma_+ + 12 a_1^6 a_1^{*2} f^2 f^{*2} g'^* \Sigma_+ + 8 a_1^4 a_1^{*4} f^2 f^{*2} g'^* \Sigma_+ + 12 a_1^6 a_1^{*2} f f^* g' g'^* \Sigma_+ + 16 a_1^8 a_1^2 f f^* g'^* \Sigma_+ + 4 a_1^{*10} g'^* \Sigma_+ + 24 a_1^4 a_1^{*6} d^* f^2 f^* \Sigma_- \Sigma_+ + 8 a_1^4 a_1^{*6} d^* f f^{*2} \Sigma_- \Sigma_+ + 16 a_1^4 a_1^{*6} d f f^* g' \Sigma_- \Sigma_+ + 16 a_1^4 a_1^{*6} d f^{*2} g' \Sigma_- \Sigma_+ + 24 a_1^2 a_1^{*8} d^* f g'^2 \Sigma_- \Sigma_+ + 6 a_1^2 a_1^{*8} d^* f^* g'^2 \Sigma_- \Sigma_+ + 2 a_1^2 a_1^{*8} d g'^3 \Sigma_- \Sigma_+ + 24 a_1^6 a_1^{*2} d f f^* g'^* \Sigma_- \Sigma_+ + 24 a_1^6 a_1^{*4} d f^{*2} g'^* \Sigma_- \Sigma_+ + 32 a_1^4 a_1^{*6} d^* f g' g'^* \Sigma_- \Sigma_+ + 8 a_1^4 a_1^{*6} d^* f^* g'^* \Sigma_- \Sigma_+ + 4 a_1^4 a_1^{*6} d g'^2 g'^* \Sigma_- \Sigma_+ + 24 a_1^6 a_1^{*4} d^* f g'^* \Sigma_- \Sigma_+ + 6 a_1^6 a_1^{*4} d^* f^* g'^* \Sigma_- \Sigma_+ + 6 a_1^6 a_1^{*4} d g' g'^* \Sigma_- \Sigma_+ + 8 a_1^8 a_1^2 d g'^* \Sigma_- \Sigma_+ + 12 a_1^4 a_1^{*6} d d^* f \Sigma_-^2 \Sigma_+ + 28 a_1^4 a_1^{*6} d d^* f^* \Sigma_-^2 \Sigma_+ + 2 a_1^4 a_1^{*6} d^2 g' \Sigma_-^2 \Sigma_+ + 12 a_1^2 a_1^{*8} d^{*2} g' \Sigma_-^2 \Sigma_+ + 3 a_1^6 a_1^{*4} d^2 g'^* \Sigma_-^2 \Sigma_+ + 8 a_1^4 a_1^{*6} d^{*2} g'^* \Sigma_-^2 \Sigma_+ + 6 a_1^6 a_1^{*4} d f^2 f^* \Sigma_+^2 + 12 a_1^6 a_1^{*4} d f f^{*2} \Sigma_+^2 + 8 a_1^4 a_1^{*6} d^* f^2 g' \Sigma_+^2 + 4 a_1^4 a_1^{*6} d^* f f^* g' \Sigma_+^2 + 2 a_1^4 a_1^{*6} d f g'^2 \Sigma_+^2 + 4 a_1^4 a_1^{*6} d f^* g'^2 \Sigma_+^2 + 12 a_1^6 a_1^{*4} d^* f^2 g'^* \Sigma_+^2 + 6 a_1^6 a_1^{*4} d^* f f^* g'^* \Sigma_+^2 + 6 a_1^6 a_1^{*4} d f g' g'^* \Sigma_+^2 + 12 a_1^6 a_1^{*4} d f^* g' g'^* \Sigma_+^2 + 12 a_1^8 a_1^2 d f g'^* \Sigma_+^2 + 24 a_1^8 a_1^2 d f^* g'^* \Sigma_+^2 + 3 a_1^6 a_1^{*4} d^2 f \Sigma_- \Sigma_+^2 + 8 a_1^4 a_1^{*6} d^{*2} f \Sigma_- \Sigma_+^2 + 12 a_1^6 a_1^{*4} d^2 f^* \Sigma_- \Sigma_+^2 + a_1^4 a_1^{*6} d^{*2} f^* \Sigma_- \Sigma_+^2 + 18 a_1^4 a_1^{*6} d d^* g' \Sigma_- \Sigma_+^2 + 27 a_1^6 a_1^{*4} d d^* g'^* \Sigma_- \Sigma_+^2 + 9 a_1^6 a_1^{*4} d d^* f \Sigma_+^3 + 2 a_1^6 a_1^{*4} d d^* f^* \Sigma_+^3 + 2 a_1^6 a_1^{*4} d^2 g' \Sigma_+^3 + 8 a_1^8 a_1^2 d^2 g'^* \Sigma_+^3.$$

These polynomials were found using a system of analytical computations *Mathematica* [4].

Appendix 2:

here we provide the proofs for lemmas stated above.

1. Due to implicit function theorem, it's sufficient to prove that Jacobian of the system (6) with respect to a set of variables (a_s, a_s^*, λ) does not vanish on the straight-line string (4). Here $s > 1$. Computing the Jacobian, we have $\det J = 2\lambda(|f|^2 - |g|^2)$. Substituting the straight-line string solution (4) to the definitions (7), we have: $\lambda = |a_1|$, $f = 0$, $g = \{a_1^2$ at $s = 2$, 0 at $s > 2\}$. Therefore, $\det J = \{-2|a_1|^3 \neq 0$ at $s = 2$, 0 at $s > 2\}$.

2. Let $s = 2$. In the vicinity of straight-line string we replace $g \rightarrow a_1^2 + \epsilon g'$, $f \rightarrow \epsilon f$, $d \rightarrow \epsilon d$, $\Sigma_\pm \rightarrow \epsilon \Sigma_\pm$. Here we parameterize the deviation of coefficients from straight-line string values by small parameter ϵ and use it as correction order counter. Let's prove that

$$a_2|_{S_\pm=0} = \sum_{n \geq 1} \frac{\epsilon^n P_n a_1^2}{a_1^{*2n} a_1^{2n}}, \quad (18)$$

The first two equations in (6) with $S_\pm = 0$ can be rewritten as

$$a_2 = -\frac{\epsilon}{a_1^{*2} a_1^2} \cdot a_1^2 \cdot (\Sigma_+ + f^* a_2^* + g'^* a_2 + d^* a_2^{*2} + \frac{1}{2} d a_2 a_2^*),$$

and it's complex conjugate. Here we have collected all ϵ -containing terms at the right hand side. Introducing variable $\tilde{\epsilon} = \epsilon/(a_1^{*2} a_1^2)$ and substituting the expansion (18), we obtain

$$\begin{aligned} \sum_{n \geq 1} \tilde{\epsilon}^n P_n &= -\tilde{\epsilon} \Sigma_+ - f^* \sum_{n \geq 1} \tilde{\epsilon}^{n+1} P_n^* a_1^{*2} \\ &- g'^* \sum_{n \geq 1} \tilde{\epsilon}^{n+1} P_n a_1^2 - d^* \sum_{n, m \geq 1} \tilde{\epsilon}^{n+m+1} P_n^* P_m^* a_1^{*4} \\ &- \frac{1}{2} d \sum_{n, m \geq 1} \tilde{\epsilon}^{n+m+1} P_n P_m^* a_1^{*2} a_1^2. \end{aligned}$$

From here we have the following recurrent relations for P_n :

$$\begin{aligned} P_1 &= -\Sigma_+, \\ P_n &= -f^* a_1^{*2} P_{n-1}^* - g'^* a_1^2 P_{n-1} \\ &- d^* a_1^{*4} \sum_{1 \leq m \leq n-2} P_{n-m-1}^* P_m^* \\ &- \frac{1}{2} d a_1^{*2} a_1^2 \sum_{1 \leq m \leq n-2} P_{n-m-1} P_m^*, \quad n > 1. \end{aligned}$$

The properties of P_n with respect to R_3, D_4 -transformations follow from the transformation laws (10) for a_2 , while the other properties of P_n can be easily proven from their recurrent definition by induction.

3. Expression (12) follows immediately from lcg-parametrization (13) and definition of lcg-oscillators given by formula (5) of Part I. Constraint χ_3 generates rotation of oscillator variables in plane (a_{n1}, a_{n2}) and rotation of vectors $\vec{e}_1, \vec{e}_2$ about $\vec{e}_3$ in opposite direction, so that the vector field $\vec{a}_{n\perp}$ is preserved. $\chi_{1,2}$ generate rotations of $\vec{e}_3$ and transformations of $\vec{a}_{n\perp}$ by formula (12). In this formula the curve $\vec{Q}$ as geometrical image is preserved, only its parametrization is changed according to (13). Components of $\vec{q}$ are proportional to gauge fixing conditions (5): $q_{s1} = \text{Re } a_{s1} = (a_s + a_{-s} + a_s^* + a_{-s}^*)/4$, $q_{s2} = \text{Re } a_{s2} =$

$i(a_s + a_{-s} - a_s^* - a_{-s}^*)/4$. Evolution, described by E_0 -transformation: $\vec{a}_{n\perp} \rightarrow \vec{a}_{n\perp} e^{-in\tau}$, is equivalent to a motion of marked point O along the curve $\vec{Q}(\sigma)$, uniform with respect to parameter (13): $\vec{Q}(\sigma) \rightarrow \vec{Q}(\sigma + \tau)$, or in terms of $(\vec{q}, \vec{p})$:

$$\begin{pmatrix} \vec{q} \\ \vec{p} \end{pmatrix} \rightarrow \begin{pmatrix} \cos s\tau & \sin s\tau \\ -\sin s\tau & \cos s\tau \end{pmatrix} \begin{pmatrix} \vec{q} \\ \vec{p} \end{pmatrix}$$

vectors $(\vec{q}, \vec{p})$ move along ellipses in tangent planes to the sphere, see fig.2. In such evolution the points $\vec{q} = 0$ move along definite curves on the sphere. On such curves the vectors $\vec{q}, \vec{p}$ are linearly dependent (corresponding ellipses degenerate to segments): $\vec{q} \times \vec{p} = 0$. The vector $\vec{q} \times \vec{p} \sim i\vec{a}_{s\perp} \times \vec{a}_{s\perp}^*$ is τ -independent and always parallel to $\vec{e}_3$. Thus, orbits of $\vec{q} = 0$ are defined as zero-level curves of a single scalar time-independent function on the sphere: $F(\vec{e}_3) = (\vec{q} \times \vec{p}, \vec{e}_3) = 0$.

4. During the period $\Delta\tau = 2\pi$ all vector fields $\vec{a}_{n\perp}$ are returned to the initial state. The field $\vec{a}_{s\perp}$ at $\Delta\tau_k^+ = \pi 2k/s$, $k = 0 \dots s-1$ is returned to the initial state, and at $\Delta\tau_k^- = \pi(2k+1)/s$, $k = 0 \dots s-1$ reverses the sign: $\vec{a}_{s\perp} \rightarrow -\vec{a}_{s\perp}$. Singular points $\vec{q} = \text{Re } \vec{a}_{s\perp} = 0$ are preserved by these transformations, while the points on the lifting generally are not.

5. For straight-line string $\vec{Q}(\sigma) \sim (\cos \sigma, \sin \sigma, 0)$, substituting it to (12) we have:


$$\vec{a}_s \sim \oint d\sigma (-\sin \sigma, \cos \sigma, 0) \exp i s (\sigma - V_x \cos \sigma - V_y \sin \sigma),$$

where $\vec{V} = \vec{e}_3$. We see that this expression does not depend on V_z and gives the same vector fields in the vicinities of northern and southern poles. For small $V_{x,y}$ we have in xy-plane, omitting unessential constant factors: $\vec{a}_{1\perp} \sim (-i, 1) + o(1)$, $\vec{q}_{1\perp} \sim (0, 1) + o(1)$, $\vec{p}_{1\perp} \sim (-1, 0) + o(1)$ - no singularity; $\vec{a}_{2\perp} \sim (-V_x - iV_y, -iV_x + V_y) + o(V_{x,y})$, $\vec{q}_{2\perp} \sim (-V_x, V_y) + o(V_{x,y})$, $\vec{p}_{2\perp} \sim (-V_y, -V_x) + o(V_{x,y})$ - saddle point; at $s > 2$ $\vec{a}_{s\perp}, \vec{q}_{s\perp}, \vec{p}_{s\perp} \sim o(V_{x,y})$ - higher order singularity. The evolution consists in rotation of vector fields in xy-plane, preserving their pole singularities.

6. The first statement follows from non-degeneracy of saddle point and general transversality theorems [10]. It also can be proven by direct computation: let's consider small deviation of solution from straight-line string (4) and associated variations of the vector fields at $s = 2$: $\vec{a} = (-V_x - iV_y, -iV_x + V_y) + (\delta a_x, \delta a_y)$, $\vec{q} = (-V_x, V_y) + (\delta q_x, \delta q_y)$, $\vec{p} = (-V_y, -V_x) + (\delta p_x, \delta p_y)$. New singular point $\vec{q} = 0$ will be slightly shifted from the pole: $V_x = \delta q_x$, $V_y = -\delta q_y$ (as a result of non-degeneracy of $\partial q_{x,y}/\partial V_{x,y}$). Considering the evolution $\vec{q} \rightarrow \vec{q} \cos 2\tau + \vec{p} \sin 2\tau$, we have $V_x = V_{x0} + b \cos 4\tau + d \sin 4\tau$, $V_y = V_{y0} + b \sin 4\tau - d \cos 4\tau$, where $V_{x0} = (\delta q_x + \delta p_y)/2$, $V_{y0} = (-\delta q_y + \delta p_x)/2$, $b = (\delta q_x - \delta p_y)/2$, $d = (\delta q_y + \delta p_x)/2$, i.e. the saddle point moves along a circle near the pole. The same orbit is defined as zero level of $(\vec{q} \times \vec{p}, \vec{V}) = (V_x - V_{x0})^2 + (V_y - V_{y0})^2 - b^2 - d^2$.

7. From lemma 2 we know, that each monomial in P_n contains at least one operator $\Sigma_{(\pm)}$, additionally $n-1$ operators from the group $(\Sigma, d, f, g'$ and their conjugates) and $2(n-1)$ operators (a_1, a_1^*) . The numbers of Σ - and d -operators are related: $n(\Sigma) = n(d) + 1$. Then we substitute (7) to the given monomial, expand it to a number of (secondary) monomials and normally order each one. Let $n(d_\uparrow), n(f_\uparrow), n(g'_\uparrow), n(a_{1\uparrow})$

be the numbers of $L_0^{(2)}$ -raising operators in the secondary monomial, $n(d_\downarrow), n(f_\downarrow), n(g'_\downarrow), n(a_{1\downarrow})$ be the numbers of $L_0^{(2)}$ -lowering operators. Using the fact, that d -operators shift $L_0^{(2)}$ by 4, f, g' -operators – by 2, $a_1^{(+)}$ by 1 (see Table 1), we conclude that in the matrix $(: P_n :)_{N_1 N_2} = \langle L_0^{(2)} = N_1 | : P_n : | L_0^{(2)} = N_2 \rangle$ the given monomial creates non-zero block with offsets $\Delta_\uparrow L_0^{(2)} = 4n(d_\uparrow) + 2n(f_\uparrow) + 2n(g'_\uparrow) + n(a_{1\uparrow})$, $\Delta_\downarrow L_0^{(2)} = 4n(d_\downarrow) + 2n(f_\downarrow) + 2n(g'_\downarrow) + n(a_{1\downarrow})$:



After summation over all secondary monomials we see that the entries in the following triangular block vanish: $N_1 + N_2 < N_{max} = \Delta_\uparrow L_0^{(2)} + \Delta_\downarrow L_0^{(2)} = 4n(d) + 2n(f) + 2n(g') + n(a_1) = 2n(d) + 2n(\Sigma) - 2 + 2n(f) + 2n(g') + 2(n-1) = 4(n-1)$.

8. The implication (i) $\Leftrightarrow$ (ii) is obvious. It's sufficient to prove (ii) for one monomial.

$$\begin{aligned} \langle L_0^{(2)} \leq N | op_{i_n} \dots op_{i_2} op_{i_1} | L_0^{(2)} \leq N \rangle &= \sum_{N_n} \dots \sum_{N_2} \sum_{N_1} \\ \langle L_0^{(2)} \leq N | op_{i_n} | L_0^{(2)} = N_n \rangle \cdot \langle L_0^{(2)} = N_n | \dots | L_0^{(2)} = N_2 \rangle \cdot \\ \langle L_0^{(2)} = N_2 | op_{i_2} | L_0^{(2)} = N_1 \rangle \cdot \langle L_0^{(2)} = N_1 | op_{i_1} | L_0^{(2)} \leq N \rangle \end{aligned}$$

The summation here is performed over all values of $L_0^{(2)}$, however, due to the $L_0^{(2)}$ -normal ordering only the values $L_0^{(2)} \leq N$ contribute, i.e. the result can be written as a product of finite matrices $op_{i_n}(N) \dots op_{i_2}(N) op_{i_1}(N)$.




Fig.7. Spectrum $(L_0^{(2)}, A_3^{(2)})$.

9. The operator a_1^2 , standing on the right of each $: P_n :$ in (16), annihilates all states of Table 2, except of $|2_1 2_{-1}\rangle$. This state is transferred by a_1^2 to $|2_{-1}\rangle$. Each monomial of

$: P_n :$ contains $n(\Sigma) \geq 1$ operators $\Sigma_\pm$ and $n_\downarrow \geq 0$ $L_0^{(2)}$ -lowering operators from the group $(a_1, f_\downarrow, g'_\downarrow, d_\downarrow)$, standing on the right. Here a_1 annihilates $|2_{-1}\rangle$, other lowering operators decrease $L_0^{(2)}$ by 2, 4 and either annihilate $|2_{-1}\rangle$ or transfer it to $|0\rangle$. The operators $\Sigma_\pm$ preserve $L_0^{(2)}$ and change $A_3^{(2)}$ by ± 1 . Figure 7 represents the spectrum of $(L_0^{(2)}, A_3^{(2)})$. Because there are no states immediately above and below $L_0^{(2)} = A_3^{(2)} = 0$ and $L_0^{(2)} = 2, A_3^{(2)} = -1$ on this figure, $\Sigma_\pm$ annihilate $|0\rangle$ and $|2_{-1}\rangle$.

Appendix 3: numerical methods.

The matrix elements of elementary operators (Table 1) in $|L_0^{(2)} \leq N\rangle$ space are represented by sparse matrices, where the typical pattern of non-zero entries is shown on fig.8. The matrices possess two-level block structure, corresponding to their $(A_3^{(2)}, L_0^{(2)})$ -charge properties. Here large blocks have constant $A_3^{(2)}$ values, inside them there are smaller blocks with constant $L_0^{(2)}$. There is an approximate similarity among the blocks. Inside the blocks of the second level the matrices possess sparse structure of non-classified type ("other!" [11]).




Fig.8. Matrix representation of elementary operators $(N = 10)$.

The most intensive part of the computation is the evaluation of large polynomials of these matrices. This is done using sparse matrix algorithms [11], where non-zero elements of the matrices are stored in condensed form. For sparse matrix multiplication two algorithms of these kind are described in [11]: pattern multiplication and threshold multiplication. The algorithms compute a product $C = A \cdot B^T$, where the transposition is computationally expensive operation due to associated changes of storage scheme, and is performed once for all operators in the Table 1. The pattern multiplication algorithm is applicable in the case when the structure of the product is known in advance and has computational cost $T_1 = O(N_C b)$, where N_C is a number of non-zero entries in C , b is an average number of non-zero entries per one row in A, B ("bandwidth"). The threshold

multiplication algorithm works in the case when the structure of the product is unknown and requires $T_2 = O(d^2b)$ operations, where d is the dimension of matrices. Multiplication of dense matrices requires $T_3 = O(d^3)$ operations, thus practically $T_1 \ll T_2 \ll T_3$. We have implemented a combination of pattern and threshold multiplication algorithms, which uses known block structure of matrices and effectively takes into account the remaining sparse structure inside the blocks. Separability of the computation to the subspaces with different values (A_3, Q) provides additional optimization. Computing the products $prod_n = prod_{n-1} \cdot op_{i(n)}^T$, we initialize $prod_0$ by the projector to a given (A_{3i}, Q_i) -subspace, represented as a sparse matrix. Although the intermediate products belong to different (A_3, Q) -blocks, this initialization automatically cuts off irrelevant contributions, considerably reducing the computational time. The localization of the final result in $\langle A_{3f}, Q_f | prod | A_{3i}, Q_i \rangle$ block, defined by its charge properties, is useful as consistency test. The resulting monomials are injected to a dense matrix a_2 , used to compute a symmetric matrix P^2 by the formula (17). Its eigenvalues are then determined by Householder algorithm [11], implemented in the library of dense matrix computations *NewMat* [12]. The computation is iterated for different correction orders $n \leq nmax = 6$. The scheme of main algorithm is given on fig.9. The necessary input data are provided by the following external program modules:




Fig.9. Scheme of computation.

- *list* – computes for the given N, s the spectrum of $(L_0^{(s)}, A_3^{(s)})$ (see fig.7) and corresponding basis in the space $|L_0^{(s)} \leq N\rangle$;
- *matr* – computes the matrix elements of the elementary operators in this basis;
- *cdeg* – *Mathematica* script, which substitutes the definitions of elementary operators in the polynomial expressions of Appendix 1, expands them to normally ordered monomials and represents the result in the form of coefficient-degree lists. The number of monomials for each order $n = 1..6$ is given in the following table:

n	1	2	3	4	5	6
num. of monomials	1	4	19	82	297	922

The determined spectrum of mass $(P^2/2\pi, S = 0, Q = 0)$ as a function of the correction order n and cutoff parameter in $L_0^{(2)} \leq N$ is presented on fig.10. The spectrum is subdivided to rows with fixed n . The points in each row show, in the order up-to-down, the spectra for $N =$

12, 16, 20, 24, 28 (sector $Q = L_0^{(2)} \bmod 4 = 0$ is defined by values $N \bmod 4 = 0$). There is no N -dependency in $n = 1$ row, because corresponding correction $P_1 = -\Sigma_+$ commutes with $L_0^{(2)}$, thus non-zero entries of P^2 for this correction are located in $\langle A_3^{(2)} L_0^{(2)} | P^2 | A_3^{(2)} L_0^{(2)} \rangle$ diagonal blocks. When N is increasing, new blocks appear, which give contributions to higher regions of P^2 , leaving intact already filled eigenvalues. Starting from $n = 2$, $L_0^{(2)}$ -nondiagonal contributions appear, leading for some eigenvalues to N -dependence saturated at large N . Other eigenvalues, as explained earlier, correspond to the states annihilated by a_2 and remain to be constant, integer and degenerate. Spectra in other Q -sectors show similar behavior.




Fig.10. $P^2/2\pi$ as a function of n, N -parameters.

Time and memory requirements of the algorithm are presented by the following statistics:

N	12	16	20	24	28
<i>subdim</i>	69	258	890	2851	8567
req.memory	1.4Mb	4.4Mb	23Mb	147Mb	1.1Gb
comp.time	2 sec	44 sec	14 min	4 hours	66 hours

Here *subdim* is the dimension of largest $(A_3 = 0, Q = 0)$ block in dense matrices, which at large N mainly defines the amount of required memory. The computation is performed on HP 2GHz Linux PC.